

**PATIENT INFORMATION**

PATIENT NAME: \_\_\_\_\_ PHONE: \_\_\_\_\_

DATE OF BIRTH: \_\_\_\_\_ SEX: ☐ M ☐ F HEIGHT: \_\_\_\_\_ WEIGHT: \_\_\_\_\_ ☐ LBS ☐ KG

ALLERGIES: \_\_\_\_\_ PREFERRED CLINIC: \_\_\_\_\_

REFERRAL STATUS: ☐ NEW REFERRAL ☐ ORDER CHANGE ☐ ORDER RENEWAL**DIAGNOSIS & CLINICAL DOCUMENTATION**☐ ICD-10 CODE: \_\_\_\_\_ DESCRIPTION: \_\_\_\_\_**REQUIRED DOCUMENTATION**☐ Insurance  
Information☐ List of  
Medications☐ Tried & Failed  
Therapies☐ Most Recent History &  
Physical☐ Recent Lab Cultures**MEDICATION ORDER**☐ Orbactiv® (Oritavancin) 1200mg IV in 1000ml **D5W** over 3 hours\*Flush with **D5W** before and after infusion☐ OTHER: \_\_\_\_\_\*Use only **D5W** for dilution and flushing, as NS is not compatible with Orbactiv®☐ REFILLS: \_\_\_\_\_

Patient to be observed for 30 minutes following the any administration.

In the event of an adverse reaction occurring in the infusion clinic, utilize the Immersiv Health adverse reaction protocol.

**PRESCRIBER INFORMATION**

PROVIDER NAME: \_\_\_\_\_ NPI #: \_\_\_\_\_

EMAIL: \_\_\_\_\_ PHONE: \_\_\_\_\_ FAX: \_\_\_\_\_

ADDRESS (INCLUDE CITY, STATE, ZIP): \_\_\_\_\_

SUPERVISING  
PHYSICIAN: \_\_\_\_\_ CONTACT NAME: \_\_\_\_\_  
(IF APPLICABLE)SIGNATURE: \_\_\_\_\_ DATE: \_\_\_\_\_  
(NO STAMPS)

SUBSTITUTION PERMITTED

DISPENSE AS WRITTEN