

PATIENT INFORMATION

PATIENT NAME: _____ PHONE: _____

DATE OF BIRTH: _____ SEX: M F HEIGHT: _____ WEIGHT: _____ LBS KG

ALLERGIES: _____ PREFERRED CLINIC: _____

REFERRAL STATUS: NEW REFERRAL ORDER CHANGE ORDER RENEWAL

DIAGNOSIS & CLINICAL DOCUMENTATION

*PLEASE COMPLETE ICD-10 FOR SPECIFIC DIAGNOSIS

- | | |
|--|--|
| <input type="checkbox"/> M04.1 Periodic Fever Syndromes (FMF, HIDS/MKD, and TRAPS) | <input type="checkbox"/> M08.2 Juvenile rheumatoid arthritis with systemic onset |
| <input type="checkbox"/> M04.2 CAPS (includes FCAS and MWS) | <input type="checkbox"/> M08.9 Juvenile arthritis |
| <input type="checkbox"/> M06.1 Adult-onset Still's Disease | <input type="checkbox"/> M10. Gout flares |
| <input type="checkbox"/> ICD-10 CODE: _____ DESCRIPTION: _____ | |

REQUIRED DOCUMENTATION

- | | | | | |
|--|--|---|---|--|
| <input type="checkbox"/> Insurance Information | <input type="checkbox"/> List of Medications | <input type="checkbox"/> Tried & Failed Therapies | <input type="checkbox"/> Most Recent History & Physical | <input type="checkbox"/> Negative TB screening |
|--|--|---|---|--|

MEDICATION ORDER

For GOUT Flare

Ilaris® (canakinumab) 150mg subcutaneous injection

For Still's Disease – SJIA & AOSD 7.5kg or greater

Ilaris® (canakinumab) 4mg/kg subcutaneous injection every 4 weeks (max dose 300mg)

For PFS – CAPS (FCAS & MWS)

15kg ≤ 40kg: Ilaris® (canakinumab) 2mg/kg subcutaneous injection every 8 weeks

15kg ≤ 40kg: Ilaris® (canakinumab) 3mg/kg subcutaneous injection every 8 weeks

> 40kg: Ilaris® (canakinumab) 150mg subcutaneous injection every 8 weeks

For PFS – FMF, HIDS/MKD & TRAPS

≤ 40kg: Ilaris® (canakinumab) 2mg/kg subcutaneous injection every 4 weeks

≤ 40kg: Ilaris® (canakinumab) 4mg/kg subcutaneous injection every 4 weeks

> 40kg: Ilaris® (canakinumab) 150mg subcutaneous injection every 4 weeks

> 40kg: Ilaris® (canakinumab) 300mg subcutaneous injection every 4 weeks

REFILL X 12 MONTHS UNLESS OTHERWISE NOTED HERE: _____

Patient to be observed for 30 minutes following the first administration. Administer per protocol.
In the event of an adverse reaction occurring in the infusion clinic, utilize the Immersiv Health adverse reaction protocol.

PRESCRIBER INFORMATION

PROVIDER NAME: _____ NPI #: _____

EMAIL: _____ PHONE: _____ FAX: _____

ADDRESS (INCLUDE CITY, STATE, ZIP): _____

SUPERVISING PHYSICIAN: _____ CONTACT NAME: _____
(IF APPLICABLE)

SIGNATURE: _____ DATE: _____
(NO STAMPS)

SUBSTITUTION PERMITTED

DISPENSE AS WRITTEN